Transparency and reproducibility in artificial intelligence

https://doi.org/10.1038/s41586-020-2766-y

Received: 1 February 2020

Accepted: 10 August 2020

Published online: 14 October 2020



Check for updates

Benjamin Haibe-Kains^{1,2,3,4,5 ⋈}, George Alexandru Adam^{3,5}, Ahmed Hosny^{6,7}, Farnoosh Khodakarami^{1,2}, Massive Analysis Quality Control (MAQC) Society Board of Directors*, Levi Waldron8, Bo Wang23,5,9,10, Chris McIntosh2,5,9, Anna Goldenberg3,5,11,12, Anshul Kundaje^{13,14}, Casey S. Greene^{15,16}, Tamara Broderick¹⁷, Michael M. Hoffman^{1,2,3,5}, Jeffrey T. Leek¹⁸, Keegan Korthauer^{19,20}, Wolfgang Huber²¹, Alvis Brazma²², Joelle Pineau^{23,24}, Robert Tibshirani^{25,26}, Trevor Hastie^{25,26}, John P. A. Ioannidis^{25,26,27,28,29}, John Quackenbush^{30,31,32} & Hugo J. W. L. Aerts^{6,7,33,34}

ARISING FROM S. M. McKinney et al. Nature https://doi.org/10.1038/s41586-019-1799-6 (2020)

Breakthroughs in artificial intelligence (AI) hold enormous potential as it can automate complex tasks and go even beyond human performance. In their study, McKinney et al. 1 showed the high potential of AI for breast cancer screening. However, the lack of details of the methods and algorithm code undermines its scientific value. Here, we identify obstacles that hinder transparent and reproducible AI research as faced by McKinney et al.1, and provide solutions to these obstacles with implications for the broader field.

The work by McKinney et al.¹ demonstrates the potential of AI in medical imaging, while highlighting the challenges of making such work reproducible. The authors assert that their system improves the speed and robustness of breast cancer screening, generalizes to populations beyond those used for training, and outperforms radiologists in specific settings. Upon successful prospective clinical validation and approval by regulatory bodies, this new system holds great potential for streamlining clinical workflows, reducing false positives, and improving patient outcomes. However, the absence of sufficiently documented methods and computer code underlying the study effectively undermines its scientific value. This shortcoming limits the evidence required for others to prospectively validate and clinically implement such technologies. By identifying obstacles hindering transparent and reproducible AI research as faced by McKinney et al.¹, we provide potential solutions with implications for the broader field.

Scientific progress depends on the ability of independent researchers to scrutinize the results of a research study, to reproduce the study's main results using its materials, and to build on them in future studies (https://www.nature.com/nature-research/editorial-policies/ reporting-standards). Publication of insufficiently documented research does not meet the core requirements underlying scientific discovery^{2,3}. Merely textual descriptions of deep-learning models can hide their high level of complexity. Nuances in the computer code may have marked effects on the training and evaluation of results⁴, potentially leading to unintended consequences⁵. Therefore, transparency in the form of the actual computer code used to train a model and arrive at its final set of parameters is essential for research reproducibility. McKinney et al.1 stated that the code used for training the models has "a large number of dependencies on internal tooling, infrastructure and hardware", and claimed that the release of the code was therefore not possible. Computational reproducibility is indispensable for high-quality Al applications^{6,7}; more complex methods demand greater transparency8. In the absence of code, reproducibility falls back on replicating methods from textual description. Although, McKinney and colleagues¹ claim that all experiments and implementation details were described in sufficient detail in the supplementary methods section of their Article¹ to "support replication with non-proprietary libraries", key details about their analysis are lacking. Even with extensive description, reproducing complex computational pipelines based purely on text is a subjective and challenging task⁹.

In addition to the reproducibility challenges inherent to purely textual descriptions of methods, the description by McKinney et al. of the model development as well as data processing and training pipelines lacks crucial details. The definitions of several hyperparameters for the model's architecture (composed of three networks referred to as the breast, lesion and case models) are missing (Table 1). In their

Princess Margaret Cancer Centre, University Health Network, Toronto, Ontario, Canada, 2Department of Medical Biophysics, University of Toronto, Toronto, Ontario, Canada, 3Department of Computer Science, University of Toronto, Toronto, Ontario, Canada. 4 Ontario Institute for Cancer Research, Toronto, Ontario, Canada. 5 Vector Institute for Artificial Intelligence, Toronto, Ontario, Canada. 6 Artificial Intelligence in Medicine (AIM) Program, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA. 7 Radiation Oncology and Radiology, Dana-Farber Cancer Institute, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA. 8 Department of Epidemiology and Biostatistics and Institute for Implementation Science in Population Health, CUNY Graduate School of Public Health and Health Policy, New York, NY, USA. 9Peter Munk Cardiac Centre, University Health Network, Toronto, Ontario, Canada. 10 Department of Laboratory Medicine and Pathobiology, University of Toronto, Ontario, Canada, 11 SickKids Research Institute, Toronto, Ontario, Canada, 12 Child and Brain Development Program, CIFAR, Toronto, Ontario, Canada. 13 Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA. 14 Department of Computer Science, Stanford University, Stanford, CA, USA. 15 Dept. of Systems Pharmacology and Translational Therapeutics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA. 16 Childhood Cancer Data Lab, Alex's Lemonade Stand Foundation, Philadelphia, PA, USA. 17 Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology, Cambridge, MA, USA. 18 Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA, 19 Department of Statistics, University of British Columbia, Vancouver, British Columbia, Canada, 20 BC Children's Hospital Research Institute, Vancouver, British Columbia, Canada. ²¹European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, Germany. ²²European Molecular Biology Laboratory, European Bioinformatics Institute, EMBL-EBI, Hinxton, UK. 23McGill University, Montreal, Quebec, Canada. 24Montreal Institute for Learning Algorithms, Quebec, Canada. ²⁵Department of Statistics, Stanford University School of Humanities and Sciences, Stanford, CA, USA. ²⁶Department of Biomedical Data Science, Stanford University School of Medicine, Stanford, CA, USA. 27 Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA. 28 Meta-Research Innovation Center at Stanford (METRICS), Stanford, CA, USA. ²⁹Department of Epidemiology and Population Health, Stanford University School of Medicine, Stanford, CA, USA. ³⁰Department of Biostatistics, Harvard T.H Chan School of Public Health, Boston, MA, USA, 31 Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA, USA, 32 Department of Data Science, Dana-Farber Cancer Institute, Boston, MA, USA, 38Radiology and Nuclear Medicine, Maastricht University, Maastricht, The Netherlands. 34Cardiovascular Imaging Research Center, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA. *A list of authors and their affiliations appears at the end of the paper. Element of the pap

Table 1 | Essential hyperparameters for reproducing the study for each of the three models

	Lesion	Breast	Case
Learning rate	Missing	0.0001	Missing
Learning rate schedule	Missing	Stated	Missing
Optimizer	Stochastic gradient Adam Missing descent with momentum		Missing
Momentum	Missing	Not applicable	Not applicable
Batch size	4	Unclear	2
Epochs	Missing	120,000	Missing

publication, McKinney et al. did not disclose the settings for the augmentation pipeline; the transformations used are stochastic and can considerably affect model performance¹⁰. Details of the training pipeline were also missing. Without this key information, independent reproduction of the training pipeline is not possible.

Numerous frameworks and platforms exist to make artificial intelligence research more transparent and reproducible (Table 2). For the sharing of code, these include Bitbucket, GitHub and GitLab, among others. The many software dependencies of large-scale machine learning applications require appropriate control of the software environment, which can be achieved through package managers including Conda, as well as container and virtualization systems, including Code Ocean, Gigantum, Colaboratory and Docker, If virtualization of the McKinney et al. internal tooling proved to be difficult, they could have released the computer code and documentation. The authors could have also created small artificial examples or used small public datasets¹¹ to show how new data must be processed to train the model and generate predictions. Sharing the fitted model (architecture along with learned parameters) should be simple aside from privacy concerns that the model may reveal sensitive information about the set of patients used to train it. Nevertheless, techniques for achieving differential privacy exist to alleviate such concerns. Many platforms allow sharing of deep learning models, including TensorFlow Hub, ModelHub.ai, ModelDepot and Model Zoo with support for several frameworks such as PvTorch and Caffe, as well as the TensorFlow library used by the authors. In addition to improving accessibility $and \, transparency, such \, resources \, can \, considerably \, accelerate \, model$ development, validation and transition into production and clinical implementation.

Another crucial aspect of ensuring reproducibility lies in access to the data the models were derived from. In their study, McKinney et al. used two large datasets under license, properly disclosing this limitation in their publication. The sharing of patient health information is highly regulated owing to privacy concerns. Despite these challenges, the sharing of raw data has become more common in biomedical literature, increasing from under 1% in the early 2000s to 20% today12. However, if the data cannot be shared, the model predictions and data labels themselves should be released, allowing further statistical analyses. Above all, concerns about data privacy should not be used as a way to distract from the requirement to release code.

Although sharing of code and data are widely seen as a crucial part of scientific research, the adoption varies across fields. In fields such as genomics, complex computational pipelines and sensitive datasets have been shared for decades¹³. Guidelines related to genomic data are clear, detailed and, most importantly, enforced. It is generally accepted that all code and data are released alongside a publication. In other fields of medicine and science as a whole, this is much less common, and data and code are rarely made available. For scientific efforts in which a clinical application is envisioned and human

Table 2 | Frameworks to share code, software dependencies and deep-learning models

	LIBI
Resource	URL
Code	
BitBucket	https://bitbucket.org
GitHub	https://github.com
GitLab	https://about.gitlab.com
Software dependencies	
Conda	https://conda.io
Code Ocean	https://codeocean.com
Gigantum	https://gigantum.com
Colaboratory	https://colab.research.google.com
Deep-learning models	
TensorFlow Hub	https://www.tensorflow.org/hub
ModelHub	http://modelhub.ai
ModelDepot	https://modeldepot.io
Model Zoo	https://modelzoo.co
Deep-learning frameworks	
TensorFlow	https://www.tensorflow.org/
Caffe	https://caffe.berkeleyvision.org/
PyTorch	https://pytorch.org/

lives would be at stake, we argue that the bar of transparency should be set even higher. If a dataset cannot be shared with the entire scientific community, because of licensing or other insurmountable issues, at a minimum a mechanism should be set so that some highlytrained, independent investigators can access the data and verify the analyses.

The lack of access to code and data in prominent scientific publications may lead to unwarranted and even potentially harmful clinical trials¹⁴. These unfortunate lessons have not been lost on journal editors and their readers. Journals have an obligation to hold authors to the standards of reproducibility that benefit not only other researchers, but also the authors themselves. Making one's methods reproducible may surface biases or shortcomings to authors before publication⁵. Preventing external validation of a model will likely reduce its impact. as it also prevents other researchers from using and building upon it in future studies. The failure of McKinnev et al. to share key materials and information transforms their work from a scientific publication open to verification and adoption by the scientific community into a promotion of a closed technology.

We have high hopes for the utility of AI methods in medicine. Ensuring that these methods meet their potential, however, requires that these studies be scientifically reproducible. The recent advances in computational virtualization and AI frameworks are greatly facilitating the implementations of complex deep neural networks in a more structured, transparent, and reproducible way. Adoption of these technologies will increase the impact of published deep-learning algorithms and accelerate the translation of these methods into clinical settings.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this paper.

Data availability

No data have been generated as part of this manuscript.

Matters arising

- McKinney, S. M. et al. International evaluation of an AI system for breast cancer screening. Nature 577, 89-94 (2020).
- Bluemke, D. A. et al. Assessing radiology research on artificial intelligence: a brief guide for authors, reviewers, and readers—from the Radiology editorial board. Radiology 293,
- Gundersen, O. E., Gil, Y. & Aha, D. W. On reproducible AI: towards reproducible research, open science, and digital scholarship in Al publications. Al Mag. 39, 56-68 (2018).
- Crane, M. Questionable answers in question answering research: reproducibility and variability of published results. Trans. Assoc. Comput. Linguist. 6, 241-252 (2018).
- Sculley, D. et al. in Advances in Neural Information Processing Systems 28 (eds Cortes, C. et al.) 2503-2511 (Curran Associates, Inc., 2015).
- Stodden, V. et al. Enhancing reproducibility for computational methods. Science 354, 1240-1241 (2016).
- Hutson, M. Artificial intelligence faces reproducibility crisis. Science 359, 725-726 (2018). Bzdok, D. & Joannidis, J. P. A. Exploration, inference, and prediction in neuroscience and
- 8. biomedicine. Trends Neurosci. 42, 251-262 (2019).
- Gundersen, O. E. & Kiensmo, S. State of the art: Reproducibility in artificial intelligence. In 9. Thirty-second AAAI Conference on Artificial Intelligence (AAAI-18) 1644–1651 (2018).
- 10. Shorten, C. & Khoshgoftaar, T. M. A survey on Image Data Augmentation for Deep Learning. J. Big Data 6, 60 (2019).
- 11. Lee, R. S. et al. A curated mammography data set for use in computer-aided detection and diagnosis research. Sci. Data 4, 170177 (2017).
- Wallach, J. D., Boyack, K. W. & Ioannidis, J. P. A. Reproducible research practices, transparency, and open access data in the biomedical literature, 2015-2017. PLoS Biol. 16, e2006930 (2018).
- 13. Amann, R. I. et al. Toward unrestricted use of public genomic data. Science 363, 350-352 (2019)
- Carlson, B. Putting oncology patients at risk. Biotechnol. Healthc. 9, 17-21 (2012).

Acknowledgements We thank S. McKinney and colleagues for their prompt and open communication regarding the materials and methods of their study. This work was supported in part by the National Cancer Institute (RO1 CA237170).

Author contributions B.H.-K. and G.A.A. wrote the first draft of the manuscript. B.H.-K. and H.J.W.L.A. designed and supervised the study. A.H., F.K., T.S., R.K., S.-A.S., W.T., R.D.W.,

C.E.M., W.J., J.D., C.F., L.W., B.W., C. McIntosh, A.G., A.K., C.S.G., T.B., M.M.H., J.T.L., K.K., W.H., A.B., J.P., R.T., T.H., J.P.A.I. and J.Q. contributed to the writing of the manuscript.

Competing interests A.H. is a shareholder of and receives consulting fees from Altis Labs. M.M.H. received a GPU Grant from Nyidia, H.J.W.L.A. is a shareholder of and receives consulting fees from Onc.Al. B.H.K. is a scientific advisor for Altis Labs. C.M. holds an equity position in Bridge7Oncology and receives royalties from RaySearch Laboratories. A.K. is on the SAB of ImmuneAl Inc, a consultant for Biogen Inc., a scientific co-founder of RavelBio Inc. and a shareholder of Freenome Inc. G.A.A., F.K., L.W., B.W., C.S.G., J.T.L., W.H., A.B., J.P., R.T., T.H., J.P.A.I. and J.Q. declare no other competing interests related to the manuscript.

Additional information

Supplementary information is available for this paper at https://doi.org/10.1038/s41586-020-

Correspondence and requests for materials should be addressed to B.H.-K. Reprints and permissions information is available at http://www.nature.com/reprints. Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2020

Massive Analysis Quality Control (MAQC) Society Board of Directors

Thakkar Shraddha³⁵, Rebecca Kusko³⁶, Susanna-Assunta Sansone³⁷, Weida Tong³⁵ Russ D. Wolfinger³⁸, Christopher E. Mason³⁹, Wendell Jones⁴⁰, Joaquin Dopazo⁴¹ & Cesare Furlanello4

³⁵National Center for Toxicological Research, US Food and Drug Administration, Jefferson, AR, USA. ³⁶Immuneering Corporation, Cambridge, MA, USA. ³⁷Engineering Science Department, Oxford e-Research Centre, University of Oxford, Oxford, UK. 38SAS Institute Inc, Cary, NC, USA. 39 Weill Cornell Medicine, New York, NY, USA. 40 Q2 Solutions, Morrisville, NC, USA. ⁴¹Hospital Virgen del Rocio, Sevilla, Spain. ⁴²Fondazione Bruno Kessler, Trento, Italy,

nature research

Corresponding author(s):

Double-blind peer review submissions: write DBPR and your manuscript number here

: instead of author names.

Last updated by author(s): YYYY-MM-DD

Reporting Summary

Nature Research wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Research policies, see our Editorial Policies and the Editorial Policy Checklist.

Statistics					
For all statistical a	nalyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.				
n/a Confirmed					
The exact	t sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement				
🗶 🗌 A statem	ent on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly				
Y	The statistical test(s) used AND whether they are one- or two-sided Only common tests should be described solely by name; describe more complex techniques in the Methods section.				
A descrip	A description of all covariates tested				
A descrip	🗷 🔲 A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons				
A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)					
	hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted uses as exact values whenever suitable.				
For Bayes	sian analysis, information on the choice of priors and Markov chain Monte Carlo settings				
For hiera	rchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes				
Estimate:	s of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated				
•	Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.				
Software ar	nd code				
Policy information	about <u>availability of computer code</u>				
Data collection	No data have been generated as part of this manuscript.				
Data analysis	No computer code has been generated as part of this manuscript.				
	g custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and encourage code deposition in a community repository (e.g. GitHub). See the Nature Research guidelines for submitting code & software for further information.				
Data					
All manuscripts n	about <u>availability of data</u> nust include a <u>data availability statement</u> . This statement should provide the following information, where applicable: es, unique identifiers, or web links for publicly available datasets				

A list of figures that have associated raw data
 A description of any restrictions on data availability
 No data have been generated as part of this manuscript.

Life sciences study design

All studies must d	isclose on these points even when	the disclosure is negative.		
Sample size	n/a			
Data exclusions	n/a			
Replication	n/a			
Randomization	n/a			
Blinding	n/a			
Reporting for specific materials, systems and methods We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response. Materials & experimental systems Methods				
n/a Involved in t	the study	n/a Involved in the study		
X Antibodie	es	ChiP-seq		
Eukaryotic cell lines		Flow cytometry		
✗ ☐ Palaeont	ology and archaeology	MRI-based neuroimaging		
X Animals a	and other organisms	·		
Human re	esearch participants			
Clinical d	ata			
Nual use	Dual use research of concern			

Reply to: Transparency and reproducibility in artificial intelligence

https://doi.org/10.1038/s41586-020-2767-x

Published online: 14 October 2020

Check for updates

Scott Mayer McKinney^{1⊠}, Alan Karthikesalingam², Daniel Tse¹, Christopher J. Kelly², Yun Liu¹, Greg S. Corrado¹ & Shravya Shetty^{1⊠}

REPLYING TO B. Haibe-Kains et al. Nature https://doi.org/10.1038/s41586-020-2766-y (2020)

We thank the authors of the accompanying Comment¹ for their interest in our work² and their thoughtful contribution. We agree that transparency and reproducibility are paramount for scientific progress. In keeping with this principle, the largest data source used in our publication is available to the academic community. Any researcher can apply for access to the OPTIMAM database (https://medphys.royalsurrey.nhs. uk/omidb/getting-access), which our institution helped fund. The broad accessibility of the database was part of the reason we pursued this collaboration. In fact, since our article came out, another group has already published results on this very dataset³.

The other dataset, from the United States, was shared with our research team after approval from the hospital system's Institutional Review Board (IRB). The IRB judged that the potential benefits of the research outweighed the minimal privacy risks associated with sharing de-identified data with a trusted party capable of and committed to safeguarding these data. As the authors understand, we are not at liberty to share data that we do not own. More generally, widely releasing data considerably alters the risk-benefit calculus for patients, so institutions must be thoughtful about how and when they do this. Because of these considerations, large medical image datasets with associated breast cancer outcomes are rarely made openly available³⁻⁵. However, as our support for the OPTIMAM database demonstrates, we endorse such efforts where practical. Although there are some small, publicly available mammography datasets⁶, restricting published research to such datasets would provide an extremely limited picture of an algorithm's clinical applicability.

The commenters¹ asked for more information concerning the training of our deep learning models. We strove to document all relevant machine learning methods while keeping the paper accessible to a clinical and general scientific audience. We thank the authors for highlighting the omission of some hyperparameters. We have supplied the requested methodological details and further elaborated on our data augmentation strategies in an Addendum⁷ to our original Article².

The authors of the Comment¹ suggest open-sourcing all the code associated with this project. Most of our work builds on open-source implementations, such as ResNet (https://github.com/tensorflow/ models/blob/master/research/slim/nets/resnet_v1.py), MobileNet (https://github.com/tensorflow/models/blob/master/research/slim/ nets/mobilenet/mobilenet v2.py), multidimensional image augmentation (https://github.com/deepmind/multidim-image-augmentation), and the Tensorflow Object Detection API (https://github.com/tensorflow/models/tree/master/research/object_detection), all of which were released by our institution. Much of the remaining code concerns data input-output and the orchestration of the training process across internal compute clusters, both of which are of scant scientific value and limited utility to researchers outside our organization. Given the extensive textual description in the supplementary information of our Article², we believe that investigators proficient in deep learning should be able to learn from and expand upon our approach.

The authors¹ further suggest releasing a containerized version of our model for others to apply to new images. It is important to note that regulators commonly classify technologies such as the one proposed here as 'medical device software' or 'software as a medical device'. Unfortunately, the release of any medical device without appropriate regulatory oversight could lead to its misuse. As such, doing so would overlook material ethical concerns. Because liability issues surrounding artificial intelligence in healthcare remain unresolved⁸, providing unrestricted access to such technologies may place patients, providers, and developers at risk. In addition, the development of impactful medical technologies must remain a sustainable venture to promote a vibrant ecosystem that supports future innovation. Parallels to hardware medical devices and pharmaceuticals may be useful to consider in this regard. Finally, increasing evidence suggests that a model's learned parameters may inadvertently expose properties of its training set to attack; how to safeguard potentially susceptible models is the subject of active research⁹. As our training data are private or under restricted access, sharing the model openly seems premature and may introduce risks that are not well characterized. On the basis of these concerns, we deliberately approach sharing artefacts derived from patient data (even if de-identified) with an abundance of caution.

No doubt the commenters¹ are motivated by protecting future patients as much as scientific principle. We share that sentiment. This work serves as an initial proof of concept, and is by no means the end of the story. We intend to subject our software to extensive testing before its use in a clinical environment, working alongside patients, providers and regulators to ensure efficacy and safety.

- Haibe-Kains, B. et al. Transparency and reproducibility in artificial intelligence. Nature https://doi.org/10.1038/s41586-020-2766-v (2020)
- McKinney, S. M. et al. International evaluation of an AI system for breast cancer screening. Nature 577, 89-94 (2020).
- Kim, H.-E. et al. Changes in cancer detection and false-positive recall in mammography using artificial intelligence: a retrospective, multireader study. Lancet Digital Health 2, e138-e148 (2020).
- Wu, N. et al. Deep neural networks improve radiologists' performance in breast cancer screening. IEEE Trans. Med. Imaging 39, 1184-1194 (2019).
- Rodriguez-Ruiz, A. et al. Stand-alone artificial intelligence for breast cancer detection in mammography: comparison with 101 radiologists. J. Natl. Cancer Inst. 111, 916-922
- Lee, R. S. et al. A curated mammography data set for use in computer-aided detection and diagnosis research. Sci. Data 4, 170177 (2017).
- McKinney, S. M. et al. Addendum: International evaluation of an AI system for breast cancer screening. Nature https://doi.org/10.1038/s41586-020-2679-9 (2020).
- Price, W. N., II, Gerke, S. & Cohen, I. G. Potential liability for physicians using artificial intelligence. J. Am. Med. Assoc. 322, 1765-1766 (2019).
- Abadi, M. et al. Deep learning with differential privacy. In Proc. 2016 ACM SIGSAC Conference Computer Communications Security CCS'16 308-318 (2016).

Matters arising

Acknowledgements We thank A. Dai and E. Gabrilovich for comments.

Author contributions This Reply was prepared by a subset of the authors of the original Article in addition to Y.L., all of whom have expertise related to this exchange. S.M.M., A.K., D.T., C.J.K, Y.L., G.S.C. and S.S. wrote and revised this Reply.

Competing interests This study was funded by Google LLC. S.M.M., A.K., D.T., C.J.K, Y.L., G.S.C. and S.S. are employees of Google and own stock as part of the standard compensation package. The authors have no other competing interests to disclose.

Additional information

 $\textbf{Correspondence and requests for materials} \ \text{should be addressed to S.M.M.} \ \text{or S.S.}$ $\textbf{Reprints and permissions information} \ is \ available \ at \ http://www.nature.com/reprints.$ $\textbf{Publisher's note} \ \textbf{Springer Nature remains neutral with regard to jurisdictional claims in}$ published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2020